

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Assistant Commissioner for Patents
United States Patent and Trademark
Office
Box PCT
Washington, D.C. 20231
ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing (day/month/year) 19 July 2000 (19.07.00)	
International application No. PCT/US99/24554	Applicant's or agent's file reference 550750-161
International filing date (day/month/year) 20 October 1999 (20.10.99)	Priority date (day/month/year) 21 October 1998 (21.10.98)
Applicant BURNSIDE, Beth, A. et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:
17 May 2000 (17.05.00)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was
☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer <p style="text-align: center;">Pascal Piriou</p> Telephone No.: (41-22) 338.83.38
--	--

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

REC'D 28 DEC 2000	
WIPO	PCT

Applicant's or agent's file reference 550750-161	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/US99/24554	International filing date (day/month/year) 20 October 1999 (20.10.1999)	Priority date (day/month/year) 21 October 1998 (21.10.1998)
International Patent Classification (IPC) or national classification and IPC IPC (7) A61K 9/16, 9/24, 9/58, 9/62 and US Cl.: 424/494, 497, 461, 462, 470		
Applicant SHIRE LABORATORIES, INC.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.

2. This REPORT consists of a total of 4 sheets, including this cover sheet.

☒ This report is also accompanied by ANNEXES, i.e., sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of 0 sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of report with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand 17 May 2000 (17.05.2000)	Date of completion of this report 11 December 2000 (11.12.2000)
Name and mailing address of the IPEA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703)305-3230	Authorized officer Alysia Berman Telephone No. 703-308-0196 DELLA MAE COLLINS PARALEGAL SPECIALIST TECHNOLOGY CENTER 1800

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International Application No.

PCT/US99/24554

I. Basis of the report

1. With regard to the elements of the international application:*

- ☒ the international application as originally filed.
- ☒ the description:
pages 1-21 _____ as originally filed
pages NONE _____, filed with the demand
pages NONE _____, filed with the letter of _____
- ☒ the claims:
pages 22-24 _____, as originally filed
pages NONE _____, as amended (together with any statement) under Article 19
pages NONE _____, filed with the demand
pages NONE _____, filed with the letter of _____
- ☒ the drawings:
pages 1-7 _____, as originally filed
pages NONE _____, filed with the demand
pages NONE _____, filed with the letter of _____
- ☒ the sequence listing part of the description:
pages NONE _____, as originally filed
pages NONE _____, filed with the demand
pages NONE _____, filed with the letter of _____

2. With regard to the language, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language _____ which is:

- ☐ the language of a translation furnished for the purposes of international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of the translation furnished for the purposes of international preliminary examination (under Rules 55.2 and/or 55.3).

3. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☐ contained in the international application in printed form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. ☒ The amendments have resulted in the cancellation of:

- ☒ the description, pages NONE
- ☒ the claims, Nos. NONE
- ☒ the drawings, sheets/fig NONE

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed, as indicated in the Supplemental Box (Rule 70.2(c)).**

* Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17).

** Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International Application No.

PCT/US99/24554

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement**1. STATEMENT**

Novelty (N)	Claims <u>1-18</u>	YES
	Claims <u>NONE</u>	NO
Inventive Step (IS)	Claims <u>6-14</u>	YES
	Claims <u>1-5, 15-18</u>	NO
Industrial Applicability (IA)	Claims <u>1-18</u>	YES
	Claims <u>NONE</u>	NO

2. CITATIONS AND EXPLANATIONS (Rule 70.7)

Claims 1-5 and 15-18 meet the criteria set out in PCT Article 33(2) and 33(4) because the prior art does not teach or fairly suggest a composition comprising one or more amphetamine salts covered with an immediate release coating and one or more amphetamine salts covered with an enteric release coating wherein the enteric release coating has a thickness of at least 25 microns.

Claims 6-14 meet the criteria set out in PCT Article 33(2)-(4), because the prior art does not teach or fairly suggest a protective layer between the drug and the enteric coating or over the enteric coating of a composition comprising immediate release coated amphetamine salts and enteric amphetamine salts.

Claims 1-5 and 15-18 lack an inventive step under PCT Article 33(3) as being obvious over OKADA et al.

OKADA et al. discloses a controlled-release pharmaceutical preparation that contains phenylpropanolamine hydrochloride in delayed pulsed release particles coated with an enteric polymer. OKADA et al. teaches that the lag time and release rate of the drug can be controlled by adding a water-soluble polymer and that the controlled release particles may be combined with immediate release preparations. The pulsed dosage form provides an equivalent blood concentration to that of the administration of several immediate release dosage forms. Figures 4 and 7 show the delayed release of the drug after 4-6 hours with subsequent rapid release of the drug. OKADA et al. teaches that the rate of release of the drug can be controlled by the thickness of the coating. It is within the skill in the art to provide the drug on or in the core. Because OKADA et al. is silent as to whether the immediate release preparations and the controlled release preparations are on the same core or different cores, this is not considered critical to the invention. It would have been obvious to one of ordinary skill in the art at the time of the invention to prepare the controlled pulsed release preparation of OKADA et al. and combine it with an immediate release coated preparation in order to increase the bioavailability of drugs.

Claims 1-18 meet the criteria set out in PCT Article 33(4) because the compositions are useful in the pharmaceutical arts.

----- NEW CITATIONS -----

VIII. Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the questions whether the claims are fully supported by the description, are made:

The description is objected to under PCT Rule 66.2(a)(v) as lacking clarity under PCT Article 5 because it fails to contain an adequate written description of the immediate release coating. The description is inadequate because: there is no mention of an immediate release coating in the specification. Therefore, one cannot determine what is encompassed by the immediate release coating.

Claims 1-18 are objected to as lacking clarity under PCT Rule 66.2(a)(v) because practice of the claimed invention is not adequately described in writing, as required under PCT Rule 5.1(a)(iii), for the reasons set forth in the immediately preceding paragraph.

Claims 1-18 are objected to under PCT Rule 66.2(a)(v) as lacking clarity under PCT Article 6 because claims 1-18 are indefinite for the following reason(s): there is no description or explanation of the immediate release coating. Therefore, the metes and bounds of the claims cannot be determined.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/24554

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 9/16, 9/24, 9/58, 9/62

US CL : 424/494, 497, 461, 462, 470

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/494, 497, 461, 462, 470

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
WEST 1.1, APS, STN: USPATFULL

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, Y	US 5837284 A (MEHTA et al.) 17 November 1998, claims 1, 2 and 23, col. 3, lines 7-10, col. 5, lines 31-36, col. 9, lines 35-40, col. 10, lines 38-42.	1-18

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

Date of mailing of the international search report

14 FEB 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Thurman K. Page

Telephone No. 703/308-1234

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/16, 9/24, 9/58, 9/62	A1	(11) International Publication Number: WO 00/23055 (43) International Publication Date: 27 April 2000 (27.04.00)
--	----	--

(21) International Application Number: PCT/US99/24554

(22) International Filing Date: 20 October 1999 (20.10.99)

(30) Priority Data:
09/176,542 21 October 1998 (21.10.98) US(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application
US 09/176,542 (CIP)
Filed on 21 October 1998 (21.10.98)

(71) Applicant (for all designated States except US): SHIRE LABORATORIES, INC. [US/US]; 1550 East Gude Drive, Rockville, MD 20850 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BURNSIDE, Beth, A. [US/US]; 1808 Briggs Chaney Road, Silver Spring, MD 20905 (US). GUO, Xiaodi [CN/US]; 7201 Dubuque Court, Derwood, MD 20855 (US). FISKE, Kimberly [US/US]; 7479 Hickory Log Circle, Columbia, MD 21045 (US). COUCH, Richard, A. [US/US]; 7620 Old Georgetown Road, Apartment 1021, Bethesda, MD 20814 (US). TREACY, Donald, J. [US/US]; 1335 Farley Court, Arnold, MD


21012 (US). CHANG, Rong-Kun [US/US]; 306 Springbrook Court, Hockessin, DE 19707 (US). MCGUINNESS, Charlotte, M. [US/US]; 6510 Broxborn Drive, Bethesda, MD 20817 (US). RUDNIC, Edward, M. [US/US]; 15103 Gravenstein Way, North Potomac, MD 20878 (US).

(74) Agents: LILLIE, Raymond et al.; Carella, Byrne, Bain, Gilfillan, Cecchi, Stewart & Olstein, 6 Becker Farm Road, Roseland, NJ 07068 (US).

(81) Designated States: AU, CA, JP, MX, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

Published*With international search report.**Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.*

(54) Title: ORAL PULSED DOSE DRUG DELIVERY SYSTEM



(57) Abstract

A multiple pulsed dose drug delivery system for pharmaceutically active amphetamine salts, comprising an immediate-release component and an enteric delayed-release component wherein (1) the enteric release coating has a defined minimum thickness and/or (2) there is a protective layer between the pharmaceutically active amphetamine salt and the enteric release coating and/or (3) there is a protective layer over the enteric release coating. The product can be composed of either one or a number of beads in a dosage form, including either capsule, tablet, or sachet method for administering the beads.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

ORAL PULSED DOSE DRUG DELIVERY SYSTEM

391 of PCT /US99/24554 filed 10-20-99, which is

This application is a continuation-in-part of Application Serial No. 09/176,542, filed October 21, 1998, the contents of which are incorporated herein by reference.

now US Patent No. 6,322,819

This invention pertains to a multiple unit dosage form delivery system comprising one or more amphetamine salts for administering the amphetamine salts to a recipient.

Background of the Invention

Traditionally, drug delivery systems have focused on constant/sustained drug output with the objective of minimizing peaks and valleys of drug concentrations in the body to optimize drug efficacy and to reduce adverse effects. A reduced dosing frequency and improved patient compliance can also be expected for the controlled/sustained release drug delivery systems, compared to immediate release preparations. However, for certain drugs, sustained release delivery is not suitable and is affected by the following factors:

First pass metabolism: Some drugs, such as β blockers, β -estradiol, and salicylamide, undergo extensive first pass metabolism and require fast drug input to saturate metabolizing enzymes in order to minimize pre-systemic metabolism. Thus, a constant/sustained oral method of delivery would result in reduced oral bioavailability.

Biological tolerance: Continuous release drug plasma profiles are often accompanied by a decline in the pharmacotherapeutic effect of the drug, e.g., biological tolerance of transdermal nitroglycerin.

Chronopharmacology and circadian rhythms: Circadian rhythms in certain physiological functions are well established. It has been recognized that many

symptoms and onset of disease occur during specific time periods of the 24 hour day, *e.g.*, asthma and angina pectoris attacks are most frequently in the morning hours (1,2).

Local therapeutic need: For the treatment of local disorders such as inflammatory bowel disease, the delivery of compounds to the site of inflammation with no loss due to absorption in the small intestine is highly desirable to achieve the therapeutic effect and to minimize side effects.

Gastric irritation or drug instability in gastric fluid: For compounds with gastric irritation or chemical instability in gastric fluid, the use of a sustained release preparation may exacerbate gastric irritation and chemical instability in gastric fluid.

Drug absorption differences in various gastrointestinal segments: In general, drug absorption is moderately slow in the stomach, rapid in the small intestine, and sharply declining in the large intestine. Compensation for changing absorption characteristics in the gastrointestinal tract may be important for some drugs. For example, it is rational for a delivery system to pump out the drug much faster when the system reaches the distal segment of the intestine, to avoid the entombment of the drug in the feces.

Pulsed dose delivery systems, prepared as either single unit or multiple unit formulations, and which are capable of releasing the drug after a predetermined time, have been studied to address the aforementioned problematic areas for sustained release preparations. These same factors are also problematic in pulsed dose formulation development. For example, gastrointestinal transit times vary not only from patient to patient but also within patients as a result of food intake, stress, and illness; thus a single-unit pulsed-release system may give higher variability compared to a multiple unit system. Additionally, drug layering or core making for multiple unit systems is a time-consuming and hard-to-optimize process. Particularly challenging for formulation scientists has been overcoming two conflicting hurdles for pulsatile formulation development, *i.e.*, lag time and rapid release.

Various enteric materials, *e.g.*, cellulose acetate phthalate, hydroxypropyl methylcellulose phthalate, polyvinyl acetate phthalate, and the EUDRAGIT® acrylic polymers, have been used as gastroresistant, enterosoluble coatings for single drug pulse

release in the intestine (3). The enteric materials, which are soluble at higher pH values, are frequently used for colon-specific delivery systems. Due to their pH-dependent attributes and the uncertainty of gastric retention time, *in-vivo* performance as well as inter- and intra-subject variability are major issues for using enteric coated systems as a time-controlled release of drugs.

A retarding swellable hydrophilic coating has been used for oral delayed release systems (4,5). It was demonstrated that lag time was linearly correlated with coating weight gain and drug release was pH independent.

Hydroxypropyl methylcellulose barriers with erodible and/or gellable characteristics formed using press coating technology for tablet dosage forms have been described to achieve time-programmed release of drugs (6). Barrier formulation variables, such as grade of hydroxypropyl methylcellulose, water-soluble and water-insoluble excipients, significantly altered the lag time and the release rate from the center cores.

Special grades of hydroxypropyl methylcellulose, *e.g.*, METOLOSE® 60SH, 90SH (Shin-Etsu Ltd., Japan), and METHOCEL® F4M (Dow Chemical Company, USA), as a hydrophilic matrix material have been used to achieve bimodal drug release for several drugs, *i.e.*, aspirin, ibuprofen, and adinazolam (7). Bimodal release is characterized by a rapid initial release, followed by a period of constant release, and finalized by a second rapid drug release.

Tablets or capsules coated with a hydrophobic wax-surfactant layer, made from an aqueous dispersion of carnauba wax, beeswax, polyoxyethylene sorbitan monooleate, and hydroxypropyl methylcellulose have been used for rapid drug release after a predetermined lag time. For example,. However, even though a two-hour lag time was achieved for the model drug theophylline at a higher coating level (60%), three hours were required for a complete release of theophylline after the lag time. (8)

A sustained-release drug delivery system is described in U.S. Patent No. 4,871,549. When this system is placed into dissolution medium or the gastrointestinal tract, water influx and the volume expansion of the swelling agent cause the explosion of the water permeable membrane. The drug thus releases after a predetermined time period.

The OROS® push-pull system (Alza Company) has been developed for pulsatile delivery of water-soluble and water-insoluble drugs (9, 10), e.g. the OROS-CT® system and is based on the swelling properties of an osmotic core compartment which provides a pH-independent, time-controlled drug release.

The PULSINCAP® dosage form releases its drug content at either a predetermined time or at a specific site (e.g., colon) in the gastrointestinal tract (11). The drug formulation is contained within a water-insoluble capsule body and is sealed with a hydrogel plug. Upon oral administration, the capsule cap dissolves in the gastric juice and the hydrogel plug swells. At a controlled and predetermined time point, the swollen plug is ejected from the PULSINCAP® dosage form and the encapsulated drug is released. A pulsatile capsule system containing captopril with release after a nominal 5-hr period was found to perform reproducibly in dissolution and gamma scintigraphy studies. However, in the majority of subjects, no measurable amounts of the drug were observed in the blood, possibly due to instability of the drug in the distal intestine. (12)

ADDERAL® comprises a mixture of four amphetamine salts which, in combination, is indicated for treatment of Attention Deficit Hyperactivity Disorder in children from 3-10 years of age. One disadvantage of current treatment is that a tablet form is commonly used which many young children have difficulty in swallowing. Another disadvantage of current treatment is that two separate doses are administered, one in the morning and one approximately 4-6 hours later, commonly away from home under other than parental supervision. This current form of treatment, therefore, requires a second treatment which is time-consuming, inconvenient and may be problematic for those children having difficulties in swallowing tablet formulations.

SUMMARY OF THE INVENTION

Accordingly, in view of a need for successfully administering a multiple unit pulsed dose of amphetamine salts and mixtures thereof, the present invention provides an oral multiple unit pulsed dose delivery system for amphetamine salts and mixtures thereof. Figure 1 illustrates the desired target plasma level profile of the pharmaceutical active contained within the delivery system.

In accordance with a preferred embodiment of the present invention, there is provided a pharmaceutical composition for delivering one or more pharmaceutically active amphetamine salts that includes:

(a) one or more pharmaceutically active amphetamine salts that are covered with an immediate release coating, and

(b) one or more pharmaceutically active amphetamine salts that are covered with an enteric release coating wherein (1) the enteric release coating has a defined minimum thickness and/or (2) there is a protective layer between the at least one pharmaceutically active amphetamine salt and the enteric release coating and/or (3) there is a protective layer over the enteric release coating.

In one embodiment, the immediate release and enteric release portions of the composition are present on the same core.

In another embodiment, the immediate release and enteric release components are present on different cores.

It is also contemplated that the composition may include a combination of the hereinabove referred to cores (one or more cores that include both components on the same core and one or more cores that include only one of the two components on the core).

The present invention provides a composition in which there is immediate release of drug and enteric release of drug wherein the enteric release is a pulsed release and wherein the drug includes one or more amphetamine salts and mixtures thereof.

The immediate release component releases the pharmaceutical agent in a pulsed dose upon oral administration of the delivery system.

The enteric release coating layer retards or delays the release of the pharmaceutical active or drug for a specified time period ("lag time") until a predetermined time, at which time the release of the drug is rapid and complete, *i.e.*, the entire dose is released within about 30-60 minutes under predetermined environmental conditions, *i.e.* a particular location within the gastrointestinal tract.

The delay or lag time will take into consideration factors such as transit times, food effects, inflammatory bowel disease, use of antacids or other medicaments which alter the pH of the GI tract.

In a preferred embodiment, the lag time period is only time-dependent, *i.e.*, pH independent. The lag time is preferably within 4 to 6 hours after oral administration of the delivery system.

In one aspect, the present invention is directed to a composition that provides for enteric release of at least one pharmaceutically active amphetamine salt, including at least one pharmaceutically active amphetamine salt that is coated with an enteric coating wherein (1) the enteric release coating has a defined minimum thickness and/or (2) there is a protective layer between the at least one pharmaceutically active amphetamine salt and the enteric release coating and/or (3) there is a protective layer over the enteric release coating.

In attempting to provide for enteric release of an amphetamine salt, applicants found that use of an enteric release coating as generally practiced in the art did not provide effective enteric release.

Typical enteric coating levels did not meet the above requirements for the desired dosage profile of amphetamine salts. Using the typical amount of enteric coating (10-20 μ) resulted in undesired premature leakage of the drug from the delivery system into the upper gastrointestinal tract and thus no drug delivery at the desired location in the gastrointestinal tract after the appropriate lag time. Thus this coating did not meet the requirements for the drug release profile to provide full beneficial therapeutic activity at the desired time.

Surprisingly, applicants found that using a thicker application of enteric coating on the formulation allowed for the second pulsed dose to be released only and completely at the appropriate time in the desired predetermined area of the gastrointestinal tract, *i.e.*, in the intestine.

This was surprising because an increase in thickness of about 5-10 μ of enteric coatings above a minimum thickness of about 10-20 μ typically does not have a significant

effect on release of drug from within such coatings. Enteric coatings typically are pH dependent and will only dissolve/disperse when exposed to the appropriate environment. Typically, application of a thicker coating (greater than 20μ) will only marginally increase the time for complete release at the appropriate environmental condition i.e., for a brief period of time (20 minutes). Using the typical coating, applicants could not achieve the desired result - rather, the coating leaked before the predetermined time in an inappropriate environment resulting in significant loss of the therapeutic agent.

Accordingly, in one aspect, the pulsed enteric release of the amphetamine salts is accomplished by employing a certain minimum thickness of the enteric coating.

In one embodiment of the invention, the pulsed dose delivery comprises a composition which comprises one or more pharmaceutically active amphetamine salts; an enteric coating over the one or more pharmaceutically active amphetamine salts, wherein the thickness of the enteric coating layer is at least 25μ ; a further layer of one or more pharmaceutically active amphetamine salts over the enteric coating layer; and an immediate release layer coating. The thicker enteric coating surprisingly provides the required delayed immediate release of the pharmaceutically active amphetamine salt at the desired time in the desired area of the gastrointestinal tract. Figure 2 illustrates a model of this delivery system.

In this aspect, the one or more pharmaceutically active amphetamine salts can be provided within or as a part of a core seed around which the enteric coating is applied. Alternatively, a core seed can be coated with one or more layers of one or more pharmaceutically active amphetamine salts.

It has further been discovered that a delayed immediate release drug delivery can also be accomplished by coating the drug first with a protective layer prior to applying the enteric coating.

Thus, in another embodiment, the pulsed enteric release is accomplished by employing a protective layer between the drug and the enteric coating. When using a protective coating, the enteric coating may be of an increased thickness or may be of lower thickness.

Thus, in another aspect, the object of the invention is met by providing a composition comprising one or more pharmaceutically active amphetamine salts; a protective layer coating over the one or more pharmaceutically active amphetamine salt layer(s), and an enteric coating layer over the protective coating layer; a further pharmaceutically active amphetamine salt layer and an immediate release layer coating. In a preferred embodiment of this aspect, the thickness of the enteric coating is at least 25 μ , and the protective layer comprises an immediate release coating.

With respect to this embodiment of the invention, the one or more pharmaceutically active amphetamine salts can be provided within or as a part of a core seed, during the core seed manufacturing process, around which the protective coating is applied. Alternatively, a core seed can be coated with one or more layers of one or more pharmaceutically active amphetamine salts.

In another embodiment, the pulsed enteric release is accomplished by employing a protective layer over the enteric coating.

Accordingly, in this embodiment of the present invention, there is provided a pulsed dose release drug delivery system comprising one or more pharmaceutically active amphetamine salts; an enteric coating layer over the pharmaceutically active amphetamine salt layer(s); and a protective layer over the enteric coating; a second pharmaceutically active amphetamine salt layer; and an immediate release layer coating.

In one aspect of this embodiment, the protective layer is comprised of one or more components, which includes an immediate release layer and a modifying layer. The modifying layer is preferably comprised of a semi water-permeable polymer. Applicants have surprisingly found that a semi-permeable polymer coating used in combination with an immediate release layer coating provided a delayed pulsed release drug delivery profile when layered over the enteric coating.

Thus, in this embodiment, the protective layer comprises a semi-permeable polymer and an immediate release coating layer. In a preferred embodiment, the modifying layer comprises a first layer of a semi-permeable polymer which is adjacent to the enteric coating

layer and a second coating layer over the semi-permeable polymer coating layer comprising an immediate release polymer coating layer.

In one aspect of this embodiment, a semi-permeable polymer, which may comprise a low water-permeable pH-insensitive polymer, is layered onto the outer surface of the enteric layer, in order to obtain prolonged delayed release time. This semi-permeable polymer coating controls the erosion of the pH-sensitive enteric polymer in an alkaline pH environment in which a pH-sensitive polymer will dissolve rapidly. Another pH-sensitive layer may be applied onto the surface of a low water-permeability layer to further delay the release time.

In a still further aspect of the invention, in addition to a protective layer, the composition comprises an acid which is incorporated into the pharmaceutical active layer or coated onto the surface of the active layer to reduce the pH value of the environment around the enteric polymer layer. The acid layer may also be applied on the outer layer of the pH-sensitive enteric polymer layer, followed by a layer of low water-permeability polymer. The release of the active thus may be delayed and the dissolution rate may be increased in an alkaline environment.

In a further embodiment, the protective coating may be used both over the drug and over the enteric coating.

With respect to this embodiment of the invention, the one or more pharmaceutically active amphetamine salts can be provided within or as a part of a core seed, during the core seed manufacturing process, around which the enteric coating is applied. Alternatively, a core seed can be coated with one or more layers of one or more pharmaceutically active amphetamine salts.

The drug delivery system of the present invention as described herein preferably comprises one or a number of beads or beadlets in a dosage form, either capsule, tablet, sachet or other method of orally administering the beads.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates a multiple pulse drug delivery system target plasma profile of the drug delivery system of the present invention. The profile reflects an immediate-release component followed by a delayed-release component.

Figure 2 schematically illustrates the delayed-release system of the present invention.

Figure 2a graphically illustrates a pulsed dose delivery system.

Figures 2b and c graphically illustrate the drug release mechanism from the proposed delivery system.

Figure 3 is a plot of the percent drug released versus time from the drug-loaded pellets described in Example 1 which exemplifies the immediate release component of the present invention.

Figure 4 is a plot of the percent drug released versus time from the coated pellets described in Example 2 which exemplifies the immediate release component and the delayed release components of the present invention.

Figure 5 is a plot of the percent drug released versus time from the coated pellets of Example 3 which exemplifies the immediate release component and the delayed release components of the present invention.

Figure 6 illustrates the drug release profile of coated pellets described in Example 4 which exemplifies the immediate release component and the delayed release components of the present invention.

Figure 7 is a plot of a profile of plasma amphetamine concentration after administration of a composite capsule containing the immediate release pellets and delayed release pellets from Examples 1 and 2, respectively.

Figure 8 is a plot of a profile of plasma amphetamine concentration after administration of a composite capsule containing the immediate release pellets and delayed release pellets from Examples 1 and 3, respectively.

DETAILED DESCRIPTION OF THE INVENTION

The present invention comprises a core or starting seed, either prepared or commercially available product. The cores or starting seeds can be sugar spheres; spheres made from microcrystalline cellulose and any suitable drug crystals.

The materials that can be employed in making drug-containing pellets are any of those commonly used in pharmaceuticals and should be selected on the basis of compatibility with the active drug and the physicochemical properties of the pellets. The additives except active drugs are chosen below as examples:

Binders such as cellulose derivatives such as methylcellulose, hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, polyvinylpyrrolidone, polyvinylpyrrolidone/vinyl acetate copolymer and the like.

Disintegration agents such as corn starch, pregelatinized starch, cross-linked carboxymethylcellulose (AC-DI-SOL®), sodium starch glycolate (EXPLOTAB®), cross-linked polyvinylpyrrolidone (PLASDONE XL®), and any disintegration agents used in tablet preparations.

Filling agents such as lactose, calcium carbonate, calcium phosphate, calcium sulfate, microcrystalline cellulose, dextran, starches, sucrose, xylitol, lactitol, mannitol, sorbitol, sodium chloride, polyethylene glycol, and the like.

Surfactants such as sodium lauryl sulfate, sorbitan monooleate, polyoxyethylene sorbitan monooleate, bile salts, glyceryl monostearate, PLURONIC® line (BASF), and the like.

Solubilizers such as citric acid, succinic acid, fumaric acid, malic acid, tartaric acid, maleic acid, glutaric acid sodium bicarbonate and sodium carbonate and the like.

Stabilizers such as any antioxidation agents, buffers, acids, and the like, can also be utilized.

Methods of manufacturing the core include

a. Extrusion-Spheronization - Drug(s) and other additives are granulated by addition of a binder solution. The wet mass is passed through an extruder equipped with a certain size screen. The extrudates are spheronized in a marumerizer. The resulting pellets are dried and sieved for further applications.

b. High-Shear Granulation - Drug(s) and other additives are dry-mixed and then the mixture is wetted by addition of a binder solution in a high shear-granulator/mixer. The granules are kneaded after wetting by the combined actions of mixing and milling. The resulting granules or pellets are dried and sieved for further applications.

c. Solution or Suspension Layering - A drug solution or dispersion with or without a binder is sprayed onto starting seeds with a certain particle size in a fluid bed processor or other suitable equipment. The drug thus is coated on the surface of the starting seeds. The drug-loaded pellets are dried for further applications.

For purposes of the present invention, the core particles have a diameter in the range of about 50-1500 microns; preferably 100-800 microns.

These particles can then be coated in a fluidized bed apparatus with an alternating sequence of coating layers.

The core may be coated directly with a layer or layers of at least one pharmaceutically active amphetamine salts and/or the pharmaceutically active amphetamine salt may be incorporated into the core material. Pharmaceutical active amphetamine salts contemplated to be within the scope of the present invention include amphetamine base, all chemical and chiral derivatives and salts thereof; methylphenidate, all chemical and chiral derivatives and salts thereof; phenylpropanolamine and its salts; and all other compounds indicated for the treatment of attention deficit hyperactivity disorder (ADHD).

A protective layer may be added on top of the pharmaceutical active containing layer and also may be provided between active layers. A separation or protective layer may be added onto the surface of the active-loaded core, and then the enteric layer is coated thereupon. Another active layer may also be added to the enteric layer to deliver an initial dose.

A protective coating layer may be applied immediately outside the core, either a drug-containing core or a drug-layered core, by conventional coating techniques such as pan coating or fluid bed coating using solutions of polymers in water or suitable organic solvents or by using aqueous polymer dispersions. Suitable materials for the protective layer include cellulose derivatives such as hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, polyvinylpyrrolidone, polyvinylpyrrolidone/vinyl acetate copolymer, ethyl cellulose aqueous dispersions (AQUACOAT®, SURELEASE®), EUDRAGIT® RL 30D, OPADRY® and the like. The suggested coating levels are from 1 to 6%, preferably 2-4 % (w/w).

The enteric coating layer is applied onto the cores with or without seal coating by conventional coating techniques, such as pan coating or fluid bed coating using solutions of polymers in water or suitable organic solvents or by using aqueous polymer dispersions. All commercially available pH-sensitive polymers are included. The pharmaceutical active is not released in the acidic stomach environment of approximately below pH 4.5, but not limited to this value. The pharmaceutical active should become available when the pH-sensitive layer dissolves at the greater pH; after a certain delayed time; or after the unit passes through the stomach. The preferred delay time is in the range of two to six hours.

Enteric polymers include cellulose acetate phthalate, Cellulose acetate trimellitate, hydroxypropyl methylcellulose phthalate, polyvinyl acetate phthalate, carboxymethylethylcellulose, co-polymerized methacrylic acid/methacrylic acid methyl esters such as, for instance, materials known under the trade name EUDRAGIT® L12.5, L100, or EUDRAGIT® S12.5, S100 or similar compounds used to obtain enteric coatings. Aqueous colloidal polymer dispersions or re-dispersions can be also applied, e.g. EUDRAGIT® L 30D-55, EUDRAGIT® L100-55, EUDRAGIT® S100, EUDRAGIT® preparation 4110D (Rohm Pharma); AQUATERIC®, AQUACOAT® CPD 30 (FMC); KOLLICOAT MAE® 30D and 30DP (BASF); EASTACRYL® 30D (Eastman Chemical).

The enteric polymers used in this invention can be modified by mixing with other known coating products that are not pH sensitive. Examples of such coating products include the neutral methacrylic acid esters with a small portion of trimethylammonioethyl methacrylate chloride, sold currently under the trade names EUDRAGIT® RS and

EUDRAGIT® RL; a neutral ester dispersion without any functional groups, sold under the trade names EUDRAGIT® NE30D; and other pH independent coating products.

The modifying component of the protective layer used over the enteric coating can include a water penetration barrier layer (semipermeable polymer) which can be successively coated after the enteric coating to reduce the water penetration rate through the enteric coating layer and thus increase the lag time of the drug release. Sustained-release coatings commonly known to one skilled in the art can be used for this purpose by conventional coating techniques such as pan coating or fluid bed coating using solutions of polymers in water or suitable organic solvents or by using aqueous polymer dispersions. For example, the following materials can be used, but not limited to: Cellulose acetate, Cellulose acetate butyrate, Cellulose acetate propionate, Ethyl cellulose, Fatty acids and their esters, Waxes, zein, and aqueous polymer dispersions such as EUDRAGIT® RS and RL 30D, EUDRAGIT® NE 30D, AQUACOAT®, SURELEASE®, cellulose acetate latex. The combination of above polymers and hydrophilic polymers such as Hydroxyethyl cellulose, Hydroxypropyl cellulose (KLUCEL®, Hercules Corp.), Hydroxypropyl methylcellulose (METHOCEL®, Dow Chemical Corp.), Polyvinylpyrrolidone can also be used.

An overcoating layer can further optionally be applied to the composition of the present invention. OPADRY®, OPADRY II® (Colorcon) and corresponding color and colorless grades from Colorcon can be used to protect the pellets from being tacky and provide colors to the product. The suggested levels of protective or color coating are from 1 to 6%, preferably 2-3 % (w/w).

Many ingredients can be incorporated into the overcoating formula, for example to provide a quicker immediate release, such as plasticizers: acetyltriethyl citrate, triethyl citrate, acetyltributyl citrate, dibutylsebacate, triacetin, polyethylene glycols, propylene glycol and the others; lubricants: talc, colloidal silica dioxide, magnesium stearate, calcium stearate, titanium dioxide, magnesium silicate, and the like.

The composition, preferably in beadlet form, can be incorporated into hard gelatin capsules, either with additional excipients, or alone. Typical excipients to be added to a capsule formulation include, but are not limited to: fillers such as microcrystalline cellulose,

soy polysaccharides, calcium phosphate dihydrate, calcium sulfate, lactose, sucrose, sorbitol, or any other inert filler. In addition, there can be flow aids such as fumed silicon dioxide, silica gel, magnesium stearate, calcium stearate or any other material imparting flow to powders. A lubricant can further be added if necessary by using polyethylene glycol, leucine, glyceryl behenate, magnesium stearate or calcium stearate.

The composition may also be incorporated into a tablet, in particular by incorporation into a tablet matrix, which rapidly disperses the particles after ingestion. In order to incorporate these particles into such a tablet, a filler/binder must be added to a table that can accept the particles, but will not allow their destruction during the tableting process. Materials that are suitable for this purpose include, but are not limited to, microcrystalline cellulose (AVICEL®), soy polysaccharide (EMCOSOY®), pre-gelatinized starches (STARCH® 1500, NATIONAL® 1551), and polyethylene glycols (CARBOWAX®). The materials should be present in the range of 5-75% (w/w), with a preferred range of 25-50% (w/w).

In addition, disintegrants are added in order to disperse the beads once the tablet is ingested. Suitable disintegrants include, but are not limited to: cross-linked sodium carboxymethyl cellulose (AC-DI-SOL®), sodium starch glycolate (EXPLOTAB®, PRIMOJEL®), and cross-linked polyvinylpyrrolidone (Plasone-XL). These materials should be present in the rate of 3-15% (w/w), with a preferred range of 5-10% (w/w).

Lubricants are also added to assure proper tableting, and these can include, but are not limited to: magnesium stearate, calcium stearate, stearic acid, polyethylene glycol, leucine, glyceryl behanate, and hydrogenated vegetable oil. These lubricants should be present in amounts from 0.1-10% (w/w), with a preferred range of 0.3-3.0% (w/w).

Tablets are formed, for example, as follows. The particles are introduced into a blender along with AVICEL®, disintegrants and lubricant, mixed for a set number of minutes to provide a homogeneous blend which is then put in the hopper of a tablet press with which tablets are compressed. The compression force used is adequate to form a tablet; however, not sufficient to fracture the beads or coatings.

It will be appreciated that the multiple dosage form of the present invention can deliver rapid and complete dosages of pharmaceutically active amphetamine salts to achieve the desired levels of the drug in a recipient over the course of about 8 hours with a single oral administration.

In so doing, the levels of drug in blood plasma of the pharmaceutically active amphetamine salts will reach a peak fairly rapidly after about 2 hours, and after about 4 hours a second pulse dose is released, wherein a second fairly rapid additive increase of plasma drug levels occurs which slowly decreases over the course of the next 12 hours.

The following examples are presented to illustrate and do not limit the invention.

EXAMPLES

Example 1

Immediate release formulation

The following formulation was used to layer the drug onto sugar spheres. Nonpareil seeds (30/35 mesh, Paulaur Corp., NJ), 6.8 kg were put into a FLM-15 fluid bed processor with a 9" Wurster column and fluidized at 60 °C. The suspension of mixed amphetamine salts (MAS) with 1% HPMC E5 Premium (Dow Chemical) as a binder was sprayed onto the seed under suitable conditions. Almost no agglomeration and no fines were observed with a yield of at least 98%. The drug-loaded cores were used to test enteric coatings and sustained release coatings.

Table 1

Ingredients	Amount (%)
Nonpareil seed	88.00
mixed amphetamine salts	11.40
METHOCEL® E5 Premium	0.60
Water	*

*removed during processing

The drug release profile of the drug-loaded pellets of this example is shown in Figure

3.

Example 2

The following formulation was used to coat the mixed amphetamine salts loaded (MASL) pellets from Example 1 with the EUDRAGIT® L 30D-55 (Rohm Pharma, Germany) coating dispersion. 2 kg of MASL pellets were loaded into a fluid bed processor with a reduced Wurster column equipped with a precision coater (MP 2/3, Niro Inc.). The coating dispersion was prepared by dispersing Triethyl citrate, Talc and EUDRAGIT® L 30D-55 into water and mixing for at least 30 minutes. Under suitable fluidization conditions, the coating dispersion was sprayed onto the fluidized MASL pellets. The spraying was continued until the targeted coating level was achieved (20 μ). The coated pellets were dried at 30-35 °C for 5 minutes before stopping the process. The enteric coated PPA pellets were tested at different pH buffers by a USP paddle method. The drug content was analyzed using HPLC. The results showed that the enteric coating delayed the drug release from the coated pellets until after exposure to pH 6 or higher (see Table 2 below). (Reference # AR98I25-4)

Table 2

Ingredients	Amount (%)
MASL pellets	40.00
EUDRAGIT® L 30D-55	24.88
Triethyl citrate	2.52
Talc	2.60
Water	*

*removed during processing

The drug release profile of the coated pellets of this example is shown in Figure 4.

Example 3

The following formulation was used to coat the MASL pellets from Example 1 with the EUDRAGIT® 4110D (Rohm Pharma, Germany) coating dispersion. MASL pellets (2 kg) were loaded in a fluid bed processor with a reduced Wurster column (GPGC-15, Glatt). The coating dispersion was prepared by dispersing Triethyl citrate, Talc and EUDRAGIT® 4110D into water and mixing for at least 30 minutes. Under suitable fluidization conditions, the coating dispersion was sprayed onto the fluidized MASL pellets. The spraying was continued until the targeted coating level was achieved. The coated pellets were dried at 30-35 °C for 5 minutes before stopping the process. The enteric coated MASL pellets were tested using a USP paddle method at different pH buffers. The drug content was analyzed using HPLC. The enteric coating delayed the drug release for several hours from the coated

pellets until the pH value reached 6.8 or higher, as shown below in Table 3. (Reference # AR98I25-3)

Table 3

Ingredients	Amount (%)
MASL pellets	70.00
Eudragit® 4110D	26.24
Triethyl citrate	0.76
Talc	3.00
Water	*

*removed during processing

The drug release profile of coated pellets of this example is shown in Figure 5.

Example 4

The following formulation was selected to coat the enteric coated MASL pellets. Coated MASL pellets from Example 2 or coated MASL pellets from Example 3 (2 kg of either) were loaded into a fluid bed processor with a reduced Wurster column (GPGC-15, Glatt). The coating dispersion was prepared by mixing SURELEASE® (Colorcon) and water for at least 15 minutes prior to spraying. Under suitable fluidization conditions, the coating dispersion was sprayed onto the fluidized pellets. The spraying was continued until the targeted coating level was achieved. The coated pellets were coated with a thin layer of OPADRY® white (Colorcon) (2%) to prevent the tackiness of the coated pellets during storage. The coated pellets were then dried at 35-40 °C for 10 minutes before discharging from the bed. The drug dissolution from both coated pellets was performed using a USP paddle method at different pH buffers. The drug content was analyzed using HPLC. The 8% SURELEASE® coating slightly sustained the drug release from EUDRAGIT® L 30D-55 coated pellets at pH 7.5 buffer, while the SURELEASE® coating delayed the drug release up to 2 hours after the buffer switched from pH 1 to pH 7.5. (Reference ## AR98I25-1)

Table 4

Ingredients	Amount, (%)
Enteric coated MASL pellets	90.00
SURELEASE®	8.00
OPADRY® white	2.00
Water	*

*removed during processing

The drug release profile of the coated pellets from this example is shown in Figure 6.

Example 5

A pulsatile delivery system can be achieved by combining the immediate release pellets (Example 1) with delayed release pellets (Example 2 or Example 3). The immediate-release pellets equivalent to half the dose and the delayed-release pellets equivalent to half the dose are filled into a hard gelatin capsule to produce the oral pulsed dose delivery system. The delayed-release portion releases the amphetamine salts rapidly and completely, after a specified lag time. The capsule products containing immediate-release pellets and delayed-release pellets (Example 1 plus Example 2 and Example 1 plus Example 3) were tested in a crossover human study. Figures 7 and 8 show the typical profiles of plasma amphetamine concentration after administration of a composite capsule containing the immediate-release pellets and delayed-release pellets from Examples 1 and 2 (10mg dose each pellet type) and a capsule containing the pellets from immediate-release pellets and delayed-release pellets from Examples 1 and 3 (10mg dose each pellet type), respectively. The general plasma profiles are similar to the desired target plasma level profile shown in Figure 1.

It is to be understood, however, that the scope of the present invention is not to be limited to the specific embodiments described above. The invention may be practiced other than as particularly described and still be within the scope of the accompanying claims.

CITED LITERATURE

1. B.Lemmer, "Circadian Rhythms and Drug Delivery", *J. Controlled Release*, 16, 63-74 (1991)
2. B. Lemmer, "Why are so many Biological Systems Periodic?" in Pulsatile Drug Delivery: Current Applications and Future Trends, R Gurny, HE Junginger and NA Peppas, eds. (Wissenschaftliche Verlagsgesellschaft mbH Stuttgart, Germany 1993) pp.11-24
3. X. Xu and PI Lee, "Programmable Drug Delivery from an Erodible Association Polymer System", *Pharm. Res.* 10(8), 1144-1152 (1993)
4. A. Gazzaniga, ME Sangalli, and F Giodano, "Oral Chonotropic Drug Delivery Systems: Achievement of Time and/or Site Specificity", *Eur J. Pharm. Biopharm.*, 40(4), 246-250 (1994)
5. A Gazzaniga, C Busetti, L Moro, ME Sangalli and F Giordano, "Time Dependent Oral Delivery Systems for Colon Targeting", *S.T.P. Pharma Sciences* 5(1), 83-88 (1996)
6. U Conte, L Maggi, ML Torre, P Giunchedi and A Lamanna, "Press-coated Tablets for Time programmed Release of Drugs", *Biomaterials*, 14(13), 1017-1023 (1993)
7. AC Shah International Patent Application WO87/00044
8. PS Walia, P Jo Mayer Stout and R Turton, "Preliminary Evaluation of an Aqueous Wax Emulsion for Controlled Release Coating", *Pharm Dev Tech*, 3(1), 103-113 (1998)
9. F Theeuwes, "OROS® Osmotic System Development", *Drug Dev Ind Pharm* 9(7), 1331-1357 (1983)
10. F Theeuwes, "Triggered, Pulsed and Programmed Drug Delivery" in Novel Drug Delivery and its Therapeutic Application, LF Prescott and WS Nimmos. eds. (Wiley, New York, 1989) pp. 323-340
11. M McNeil, A Rashid and H Stevens, " International Patent App WO90/09168

12. IR Wilding, SS Davis, M Bakhshae, HNE Stevens, RA Sparrow and J Brennan, "Gastrointestinal Transit and Systemic Absorption of Captopril from a Pulsed Release Formulation", Pharm Res 9(5), 654-657 (1992)

CLAIMS:

1. A pharmaceutical composition for enteric delivery of one or more pharmaceutically active amphetamine salts comprising:

(a) one or more pharmaceutically active amphetamine salts covered with an immediate release coating; and

(b) one or more pharmaceutically active amphetamine salts that are covered with an enteric release coating that provides for pulsed enteric release wherein the enteric release coating has a thickness of at least 25 μ .

2. The pharmaceutical composition of claim 1 wherein the one or more pharmaceutically active amphetamine salts are coated onto a core.

3. The pharmaceutical composition of claim 1 wherein the one or more pharmaceutically active amphetamine salts are incorporated into a core.

4. The pharmaceutical composition of claim 1, wherein the immediate release and enteric release portions of the composition are present on a single core.

5. The pharmaceutical composition of claim 1, wherein the immediate release and enteric release components are present on different cores.

6. A pharmaceutical composition for enteric delivery of one or more pharmaceutically active amphetamine salts comprising:

(a) one or more pharmaceutically active amphetamine salts covered with an immediate release coating;

(b) one or more pharmaceutically active amphetamine salts that are covered with an enteric release coating that provides for pulsed enteric release; and

(c) a protective layer between the at least one pharmaceutically active amphetamine salt and the enteric release coating.

7. The pharmaceutical composition of claim 6 wherein the one or more pharmaceutically active amphetamine salts are coated onto a core.

8. The pharmaceutical composition of claim 6 wherein the one or more pharmaceutically active amphetamine salts are incorporated into a core.
9. The pharmaceutical composition of claim 6 wherein the enteric release coating has a minimum thickness of 25 μ .
10. A pharmaceutical composition for delivering one or more pharmaceutically active amphetamine salts comprising:
 - (a) one or more pharmaceutically active amphetamine salts covered with an immediate release coating;
 - (b) one or more pharmaceutically active amphetamine salts that are covered with an enteric release coating that provides for pulsed enteric release; and
 - (c) a protective layer over the enteric release coating.
11. The pharmaceutical composition of claim 10 wherein the one or more pharmaceutically active amphetamine salts are coated onto a core.
12. The pharmaceutical composition of claim 10 wherein the one or more pharmaceutically active amphetamine salts are incorporated into a core.
13. The pharmaceutical composition of claim 10, wherein the immediate release and enteric release portions of the composition are present on a single core.
14. The pharmaceutical composition of claim 10, wherein the immediate release and enteric release components are present on different cores.
15. A pharmaceutical composition for delivery of at least one amphetamine salt, comprising:
 - (a) at least one pharmaceutically active amphetamine salt covered with an immediate release coating; and
 - (b) at least one pharmaceutically active amphetamine salt covered with an enteric release coating, said component (a) providing for an immediate release of amphetamine salt to provide a first blood level of amphetamine salt and component (b)

providing a delayed pulsed-release of amphetamine salt that increases the blood level of amphetamine salt to a second level that is greater than the first level provided by component (a).

16. The pharmaceutical composition of claim 15 wherein the enteric release coating has a minimum thickness of 25 microns.

17. The pharmaceutical composition of claim 15 wherein the delayed pulsed-release is from 4 to 6 hours after administration of the pharmaceutical composition.

18. The pharmaceutical composition of claim 15 wherein the delayed pulsed-release, releases the amphetamine salt in about 30 to 60 minutes after initiation of the release.

Figure 1




FIGURE 2A




FIGURE 2B




FIGURE 2C




FIGURE 3




Figure 4




Figure 5





Figure 6







Fig. 7

*Fig. 8*

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/24554

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 9/16, 9/24, 9/58, 9/62

US CL : 424/494, 497, 461, 462, 470

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/494, 497, 461, 462, 470

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
WEST 1.1, APS, STN: USPATFULL

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, Y	US 5837284 A (MEHTA et al.) 17 November 1998, claims 1, 2 and 23, col. 3, lines 7-10, col. 5, lines 31-36, col. 9, lines 35-40, col. 10, lines 38-42.	1-18



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

Date of mailing of the international search report

14 FEB 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Thurman K. Page

Telephone No. 703/308-1234